


Probabilistic
Graphical
Models



Introduction

Preliminaries: Distributions

Joint Distribution

$P(I, D, G)$

- Intelligence (I) ← 2
 - i^0 (low), i^1 (high),
 - Difficulty (D) ← 2
 - d^0 (easy), d^1 (hard)
 - Grade (G) ← 3
 - g^1 (A), g^2 (B), g^3 (C)
- parameters
 $2 \times 2 \times 3 = 12$
- independent params
 ||

I	D	G	Prob.
i^0	d^0	g^1	0.126
i^0	d^0	g^2	0.168
i^0	d^0	g^3	0.126
i^0	d^1	g^1	0.009
i^0	d^1	g^2	0.045
i^0	d^1	g^3	0.126
i^1	d^0	g^1	0.252
i^1	d^0	g^2	0.0224
i^1	d^0	g^3	0.0056
i^1	d^1	g^1	0.06
i^1	d^1	g^2	0.036
i^1	d^1	g^3	0.024

↓

Conditioning

condition on g^1

I	D	G	Prob.
i^0	d^0	g^1	0.126
i^0	d^0	g^2	0.168
i^0	d^0	g^3	0.126
i^0	d^1	g^1	0.009
i^0	d^1	g^2	0.045
i^0	d^1	g^3	0.126
i^1	d^0	g^1	0.252
i^1	d^0	g^2	0.0224
i^1	d^0	g^3	0.0056
i^1	d^1	g^1	0.06
i^1	d^1	g^2	0.036
i^1	d^1	g^3	0.024


Conditioning: Reduction

I	D	G	Prob.
i^0	d^0	g^1	0.126
i^0	d^1	g^1	0.009
i^1	d^0	g^1	0.252
i^1	d^1	g^1	0.06

Conditioning: Renormalization

I	D	G	Prob.
i^0	d^0	g^1	0.126
i^0	d^1	g^1	0.009
i^1	d^0	g^1	0.252
i^1	d^1	g^1	0.06

$P(I, D, g^1)$
unnormalized measure 0.447



I	D	Prob.
i^0	d^0	0.282
i^0	d^1	0.02
i^1	d^0	0.564
i^1	d^1	0.134

$P(I, D | g^1)$

Marginalization

$P(I, D)$

Marginalize I

I	D	Prob.
i^0	d^0	0.282
i^0	d^1	0.02
i^1	d^0	0.564
i^1	d^1	0.134

D	Prob.
d^0	0.846
d^1	0.154